



## 醫學美容培訓班

日期:8月25 - 26日 時間:12:00 - 19:00 課程人數:30人

地點:澳門大醫院 525 室

課程完成後,學員將獲發澳門健康管理及促進學會培訓證書

- 課程是以面對面的形式進行,約14小時。
- 参加者必 須年滿 18 歲,學員需於開始上課前至少 10 分鐘到達。
  - 查詢電話 (853 2855 6207), 報名請於 8 月 23 日前把姓名、工作單位名稱及聯絡電話電郵 aizza.event.vivian@gmail.com

報名費: \$300/人 (包括培訓器材及相關餐飲) 此培訓班由 澳門基金會資助部份活動經費



支持單位



澳門科技大學健康科學學院 專業醫療發展中心



澳門美容專業協會

### 課程內容

### 8月25日:

- 1. 再生療程的發展
- 2. 雙頻點陣激光的秘密
- 3. 私密處護理的秘密
- 4. 案例分享
  - > 女性尿滲問題的處理
  - ▶ 解構產後陰道鬆弛
- 5 手把手培訓班·
  - ▶ 面部雙頻點陣激光療程
  - > 女性陰道激光療程

### 8月26日

- 6. 非入侵式、微創式醫美療程概覽
- 7. 醫學美容技術簡介:
  - ▶超聲波爆脂
  - ▶ 微針療法
- 8. 手把手培訓班:
  - ▶超聲波爆脂
  - ▶ 微針療法

授課語言:粤語和英語

## 講者/導師

潘廣嗣口腔醫院院長(中國) 陳祖鈞整形外科醫生(香港) 黃天厚皮膚學碩士(英國) 臧薇敏口腔醫生(香港) 張 晨口腔醫生(香港)


陳敏醫生(澳門) 澳門衞生局婦產科專科文憑






以前的觀念是生產後一個月內應盡可能休息,甚至不要 下來,但目前較新的觀念則認為產後愈年運動,則身體 的程度極快。 產後可做的運動一、胸部運動(養後第2天開始); 產後可做的運動一、胸部運動(產後第2天開始); 二、乳部運動(產後第3天開始); 三、乳部運動(產後第3天開始); 三、頸部運動(產後第5天開始); 五、臀部運動(產後第5天開始); 九、收輸陸部運動(產後第10天開始); 七、子宮收施運動(應後第10天開始); 七、子宮收施運動(電後第10天開始); 1、即即應動應後後第15天開始); 1、即即應動應後後第15天開始); 1、即以應動(產後第15天開始);

















Mechanism of Action — Hybrid Fractional Laser diVa distinctions — deeper ablation and coagulation

- 2940nm is used for precise ablation
  - 0 800 µm
- 1470 nm is used for coagulation of epithelium and lamina propia
  - 100 700 µm
- This unique blend provides customizable treatments for epithelial tissue to heal quickly and the lamina propria to remodel slowly over time, laying down more collagen, vascularity, and elastin.





# What is Hybrid Fractional Laser (HFL)? • 2 wavelengths 2940 Erbium (Ablation) 1470 Diode (Coagulation) • Why is this important? - CUSTOMIZATION - Different conditions need different treatments - 1st treatment settings can be different from 3rd treatment


2

























## Physician Investigator's Recommended Treatment Protocol 1. Review medical and gynecologic history 2. Confer with patient's GYN before procedure if you deem that necessary 3. Obtain signed informed consent 4. Pregnancy test (if of childbearing potential) 5. Patient should empty bladder prior to treatment 6. Apply topical numbing for 20-30 min 7. Position patient in dorsal recumbent position with buttocks at end of table and/or feet in stirrups 8. Swab off topical numbing





